

The Discrete Fractional Fourier Transform and Harper's Equation

Laurence Barker

Technical Report BU-CE-0007,
Department of Computer Engineering, Bilkent University.
Preprint April 1999, this version April 2000.

Department of Mathematics, Bilkent University, 06533 Bilkent, Ankara, Turkey. barker@fen.bilkent.edu.tr	39A12 Finite differences: Discrete version of topics in analysis. 47N40 Operator Theory: Applications to numerical analysis. 65T20 Numerical Analysis: Discrete and fast Fourier transforms.
--	---

Abstract We show that the discrete fractional Fourier transform recovers the continuum fractional Fourier transform via a limiting process whereby inner products are preserved.

1: Discussion

A generalization of the Fourier Transform (FT), the fractional FT of Namias [12] is widely used in signal-processing, optics, and quantum mechanics; see Özaktaş–Kutay–Mendlovic [13] for discussions of applications and history. The discrete FT, already a valuable tool in classical number theory during the 1930's, acquired, with the advent of computers, a tremendous range of applications throughout science and engineering; any calculation executed by a digital machine is, by nature, discrete. Pei–Yeh [15] introduced the discrete fractional FT. Candan [7] proved that the discrete fractional FT is uniquely determined by its defining properties; see Candan–Kutay–Özaktaş [8]. The eigenvectors of the discrete fractional FT, which we call the *discrete Hermite–Gaussian functions*, “converge” to the eigenvectors of the fractional FT, which are the Hermite–Gaussian functions. As operators, the discrete fractional FT “converges” to the fractional FT. Numerical evidence for these “convergences” was given in [15], and a heuristic argument was given in [7] and [8].

As often happens when rigorous justification of a belief emerges, the principal benefit may be not so much in the confirmation itself as in the precise formulation of the assertion. We shall drop the vague term “convergence”, since we shall be concerned with sequences whose terms all belong to different spaces. Rather, in Theorems 2.5 and 2.8, respectively, we say that the Harper functions *induce* the Hermite–Gaussians, and that the discrete fractional FT *induces* the fractional FT; the meaning is still that, as the underlying period increases, the discrete constructions, in some (particular) sense, approach the continuum constructions.

An important feature of both the discrete fractional FT and the continuum fractional FT is that they are unitary transformations. This being so, we must recognise that the spaces upon which they operate are inner product spaces. The notion of induction is intended to describe discrete to continuum correspondences that preserve inner products. A systematic examination of induction is given in [5]. Some applications of the induction property of the fractional FT are described in [6]. The purpose of the the present paper is to give proofs of Theorems 2.5 and 2.8, which are invoked in the other two papers. The first version (April

1999) of the present paper was composed before [5] and [6], and the discussions below are still self-contained. Actually, the author considers this to be a disadvantage; the account could be abbreviated by means of references to the two later papers. Furthermore, some of the technical arguments below are more cumbersome than need be. A more streamlined presentation of the work is in preparation.

Balian–Itzykson [4], Athanasiu–Floratos [1], Athanasiu–Floratos–Nicolis [2], Hakioglu [9] have studied discrete quantum systems that admit a symmetry under the action of a suitable cyclic subgroup of $SL_2(q)$, where q is a prime-power. Any connection with symmetries from $SL_2(\mathbb{R})$ for continuum quantum systems remains, as yet, unclear; there is little reason to expect any deep connection, since $SL_2(\mathbb{R})$ does not seem to be realizable in any way as a continuum limit of $SL_2(q)$. However, [17], [1], [2], [9] suggested to us that the FT be taken as a focus for realizing continuum quantum systems as limits of discrete quantum systems: the relevant representation of $SL_2(\mathbb{R})$ sends the matrix $(0, -1; 1, 0)$ to the FT; the relevant representation of $SL_2(q)$ sends the matrix $(0, -1; 1, 0)$ to the discrete FT.

Let us indicate the ways in which Harper’s Equation provides a rapport between the material in this paper and a couple of topics in quantum mechanics. Consider an integer $n \geq 1$, let ψ be a function $\mathbb{Z} \rightarrow \mathbb{C}$ with period dividing n , and let $E \in \mathbb{R}$. Harper’s Equation is

$$\psi(x-1) + 2\cos(2\pi x/n)\psi(x) + \psi(x+1) = (4 - 2\pi E/n)\psi(x)$$

where $x \in \mathbb{Z}$. Assume now that ψ and E are solutions to Harper’s Equation. Then ψ is a Hamiltonian eigenstate for a discrete analogue of a quantum harmonic oscillator. To see this in a quick informal way, take n to be large, write $\xi = x\sqrt{2\pi/n}$, and suppose that $\xi \ll \sqrt{n}$. Put

$$\phi(\xi) = (n/2\pi)^{1/4}\psi(x).$$

(The factor $(n/2\pi)^{1/4}$ is just for normalization, and can be ignored for the purpose of the present discussion.) Then Harper’s Equation reduces to

$$n/2\pi(-\phi(\xi - \sqrt{2\pi/n}) + 2\phi(\xi) - \phi(\xi + \sqrt{2\pi/n})) + \xi^2\phi(\xi) = E\phi(\xi) + O((\xi/n)^2).$$

Thus we have obtained an approximation to a difference analogue of the Schrödinger equation

$$\left(-\frac{d^2}{dr^2} + r^2 - E\right)\theta(r) = 0$$

of a harmonic oscillator. The Hamiltonian eigenstates of this harmonic oscillator are the solutions $\theta = h_j$, where h_j is the j -th Hermite–Gaussian function $\mathbb{R} \rightarrow \mathbb{C}$ (but with real values) given by

$$h_j(r) = \pi^{-1/4}2^{-j/2}(j!)^{-1/2}e^{-r^2}H_j(r)$$

and H_j is the Hermite polynomial of degree j given by

$$H_j(r) = (-1)^j e^{r^2} \frac{d^j}{dr^j} e^{-r^2/2}.$$

Here, the index j runs over the set $\mathbb{N} = \{0, 1, 2, \dots\}$ of natural numbers. The energy eigenvalue corresponding to the eigenstate h_j is $2j + 1$. In other words,

$$\left(-\frac{d^2}{dr^2} + r^2\right)h_j(r) = (2j + 1)h_j(r).$$

In anticipation of a comment below, we recall that the above Schrödinger equation also has a family of “physically uninteresting” solutions which behave as $e^{r^2/2}$. We also mention that the time-evolution operator of the above harmonic oscillator is a (time-dependent) scalar multiple of the fractional FT.

In Atakishiev–Suslov [3], two other discrete analogues of a harmonic oscillator are examined in connection with two other discrete analogues of the Hermite–Gaussians, namely, the Kravchuk polynomials and the Hermite q -polynomials. It is an open question as to what relationship, if any, exists between the discrete Hermite Gaussians and either the Kravchuk polynomials or the Hermite q -polynomials.

For a Bloch electron confined to the lattice $\mathbb{Z} \times \mathbb{Z} = \{(a, b) : a, b \in \mathbb{Z}\}$ and subject to a uniform transverse time-invariant magnetic field, let us impose the Landau gauge. Then the Hamiltonian \widehat{h} acts on a wave function θ such that

$$(\widehat{h}\theta)(a, b) = \theta(a - 1, b) + \theta(a + 1, b) + e^{-2\pi i \omega a} \theta(a, b - 1) + e^{2\pi i \omega a} \theta(a, b + 1)$$

where ω is a real constant proportional to the magnetic flux. To obtain periodic solutions, let us assume that ω is rational, and write $\omega = m/n$ where m and n are coprime integers, and $n \geq 1$. For the Schrödinger equation $\widehat{h}\theta = H\theta$, the n^2 solutions with period dividing n coincide with the solutions to

$$\theta(a, b) = e^{2\pi i kb/n} \phi(a)$$

where k is an integer, and

$$\phi(a - 1) + \phi(a + 1) + 2 \cos(2\pi(ma + k)/n) \phi(a) = H\phi(a).$$

This form of Harper’s Equation is essentially no more general than before. Indeed, since m is coprime to n , the change of variables $a \mapsto ma + k$ is invertible up to congruence modulo n , and we can put $\psi(ma + k) = \phi(a)$ and $E = n(4 - H)/2\pi$. For a study of Bloch electrons subjected to uniform magnetic fields see, for instance, Rammal–Bellissard [17].

Harper’s Equation (as an abstract system of linear equations) is the starting point for the Pei–Yeh definition of the discrete fractional FT. The Harper functions are defined to be precisely those solutions to Harper’s Equation which commute with the discrete FT; see Section 2. Theorem 2.5, below, may be interpreted as saying that, for fixed $F \in \mathbb{R}$, and as n increases, the Harper functions with eigenvalues at least $4 - 2\pi F/n$ induce the Hermite–Gaussians. This conclusion was perhaps to be expected in view of the physics above, and we have already indicated the numerical and heuristic evidence. Without Theorem 2.5, how could we be sure that no Harper functions with high eigenvalues induce the “physically uninteresting” solutions to the Schrödinger equation of the above harmonical oscillator? Admittedly, this particular query can easily be resolved using Lemma 3.4. But how could we be sure that, for large n , there are no Harper functions which have high eigenvalues, yet have nothing to do with the Hermite–Gaussians?

For an integer $n \geq 1$, let $L(\mathbb{Z}(n))$ denote the space of functions $\psi : \mathbb{Z} \rightarrow \mathbb{C}$ such that $\psi(x + n) = \psi(x)$ for all $x \in \mathbb{Z}$. Let $\mathbb{Z}(n)$ denote the set consisting of the integers x such that $-n/2 < x \leq n/2$. We define addition on $\mathbb{Z}(n)$ so as to respect the usual addition in \mathbb{Z} up to congruence modulo n . Thus, given $x, y, z \in \mathbb{Z}(n)$, then $x + y = z$ provided n divides $x + y - z$. We define an inner product on $L(\mathbb{Z}(n))$ by

$$\langle \phi | \psi \rangle := \sum_{x \in \mathbb{Z}(n)} \phi(x)^* \psi(x).$$

We shall establish some connections between a continuum scenario based on the complex Hilbert space $L^2(\mathbb{R})$ and discrete scenarios based on the finite-dimensional spaces $L(\mathbb{Z}(n))$ in the limit as $n \rightarrow \infty$. Some scaling factors will be involved, and it is convenient to define

$$\nu(n) := (n/2\pi)^{1/4}.$$

The period n will not be allowed to range over all of \mathbb{N} . We shall confine n to an infinite subset $\mathcal{N} \subset \mathbb{N}$ satisfying a suitable hypothesis. Vectors in $L^2(\mathbb{R})$ are to be related to some sequences of the form $(\psi_n : n \in \mathcal{N})$, where each $\psi_n \in L(\mathbb{Z}(n))$. Operators on $L^2(\mathbb{R})$ are to be related to some sequences of the form $(\hat{a}_n : n \in \mathcal{N})$, where each a_n is an operator on $L(\mathbb{Z}(n))$. Some technical care will be needed, and in order that the wood not be lost for the trees, let us indicate, in a casual way, the rough idea. Consider a vector $\psi \in L^2(\mathbb{R})$, and for simplicity of discussion, let us assume that ψ is rapidly decreasing. Loosely speaking, a sequence $(\psi_n : n \in \mathcal{N})$ with $\psi_n \in L(\mathbb{Z}(n))$ will be deemed to be related to the vector ψ provided we have an approximate equality

$$\psi(x\nu(n)^{-2}) \simeq \nu(n)\psi_n(x)$$

for large n and for most of the elements $x \in \mathbb{Z}(n)$. The analogous relationship between operators will be constructed in such a way as to respect the relationship between vectors. This idea, in some form or another, has always been used in realizations of the FT as a continuum limit of the discrete FT. In Section 2, we express this idea in an algebraic way that will suit our purposes here and in subsequent work. We also review the Pei–Yeh definition of the discrete fractional FT, and we state our main results, Theorems 2.5 and 2.8. The four Theorems stated in Section 2 are proved in Section 3.

2: Results

Recall that a smooth function $\phi : \mathbb{R} \rightarrow \mathbb{C}$ is said to be **rapidly decreasing** provided, for all $a, b \in \mathbb{N}$, the function $\mathbb{R} \rightarrow \mathbb{R}$ given by $r \mapsto |\phi^{(a)}(r)r^b|$ has an upper bound. A dense subspace of $L^2(\mathbb{R})$, the **Schwartz space** $\mathcal{S}(\mathbb{R})$ is defined to be the space of rapidly-decreasing functions. It is well-known that the set $\{h_j : j \in \mathbb{N}\}$ of Hermite-Gaussians is a complete set in $\mathcal{S}(\mathbb{R})$, and is an orthonormal basis for the Hilbert space $L^2(\mathbb{R})$. Following Namias [12], we define, for each $t \in \mathbb{R}$, an operator $\hat{f}^{[t]}$ on $L^2(\mathbb{R})$ given by

$$\hat{f}^{[t]}h_j = e^{2\pi ijt}h_j.$$

Namias showed that, for $0 < t < 1/2$ and $\phi \in \mathcal{S}(\mathbb{R})$, we have

$$(\hat{f}^{[t]}\phi)(r) = (2\pi \sin 2\pi t)^{-1/2} \int_{-\infty}^{\infty} K_t(r, q)\phi(q)dq$$

where the kernel is

$$K_t(r, q) = \exp(i\pi(t - 1/4) - i(\cot 2\pi t)(r^2 + q^2)/2 + ir/\sin 2\pi t).$$

As observed in McBride–Kerr [10], the action of $\hat{f}^{[t]}$ on $\mathcal{S}(\mathbb{R})$ for arbitrary $t \in \mathbb{R}$ is determined by the explicit integral formula together with the condition that $\hat{f}^{[t+t']} = \hat{f}^{[t]}\hat{f}^{[t']}$ for all $t, t' \in \mathbb{R}$. The Fourier Transform \hat{f} on $L^2(\mathbb{R})$ is the special case

$$\hat{f} = \hat{f}^{[1/4]}$$

so it makes sense to call $\widehat{f}^{[t]}$ the **fractional Fourier Transform**.

The operator on $\mathcal{S}(\mathbb{R})$ given by $\phi(r) \mapsto (-d^2/dr^2 + r^2)\phi(r)$ extends to an operator \widehat{h} on $L^2(\mathbb{R})$ given by

$$\widehat{h}h_j = (2j + 1)h_j$$

for $j \in \mathbb{N}$. Evidently, $\widehat{f}^{[t]} = \exp(i\pi t(\widehat{h} - 1))$.

Consider an integer $n \geq 1$. For the moment, let us take n to be fixed. Recall that the **discrete Fourier Transform** \widehat{f}_n is defined to be the operator on $L(\mathbb{Z}(n))$ such that

$$(\widehat{f}_n\psi)(x) = \frac{1}{\sqrt{n}} \sum_{y \in \mathbb{Z}(n)} e^{2\pi ixy/n} \psi(y)$$

where $\psi \in L(\mathbb{Z}(n))$, and $x \in \mathbb{Z}$. For each integer x , we write x^* to denote the function $\mathbb{Z} \rightarrow \mathbb{C}$ such that $x^*(y) = 1$ when $y \equiv x \pmod{n}$, and $x^*(y) = 0$ otherwise. The set $\{x^* : x \in \mathbb{Z}(n)\}$ is an orthonormal basis for $L(\mathbb{Z}(n))$. It is easily shown that

$$\widehat{f}_n x^* = \frac{1}{\sqrt{n}} \sum_{y \in \mathbb{Z}(n)} e^{2\pi ixy/n} y^*.$$

Another easy calculation yields $\widehat{f}_n^2 x^* = (-x)^*$, whence $\widehat{f}_n^4 = 1$.

Let \widehat{u}_n and \widehat{v}_n be the unitary operators on $L(\mathbb{Z}(n))$ such that $\widehat{u}_n x^* = (x + 1)^*$ and $\widehat{v}_n x^* = e^{2\pi ix/n} x^*$ for $x \in \mathbb{Z}$. The hermitian conjugates satisfy $\widehat{u}_n^\dagger x^* = (x - 1)^*$ and $\widehat{v}_n^\dagger x^* = e^{-2\pi ix/n} x^*$. We define a hermitian operator

$$\widehat{s}_n := \widehat{u}_n + \widehat{u}_n^\dagger + \widehat{v}_n + \widehat{v}_n^\dagger$$

on $L(\mathbb{Z}(n))$. Given an eigenvector σ of \widehat{s}_n with corresponding eigenvalue $4 - 2\pi E/n$, then σ and E satisfy Harper's Equation

$$\sigma(x - 1) + 2 \cos(2\pi x/n) \sigma(x) + \sigma(x + 1) = (4 - 2\pi E/n) \sigma(x).$$

REMARK 2.1: *Let σ be an eigenvector of \widehat{s}_n . Then there does not exist an integer x such that $\sigma(x) = 0 = \sigma(x + 1)$.*

Proof: This is immediate from the fact that σ satisfies Harper's Equation. \square

THEOREM 2.2: (Candan [7]) *Up to scalar multiplication, $L(\mathbb{Z}(n))$ has a unique basis consisting of common eigenvectors of \widehat{s}_n and \widehat{f}_n .*

Proof: We give a brief presentation of the argument in Candan–Kutay–Özaktaş [8, Section III.B]. By direct calculation, $\widehat{f}_n \widehat{u}_n \widehat{f}_n^\dagger = \widehat{v}_n$ and $\widehat{f}_n \widehat{v}_n \widehat{f}_n^\dagger = \widehat{u}_n^\dagger$. Hence $\widehat{f}_n \widehat{s}_n = \widehat{s}_n \widehat{f}_n$. In particular, $L(\mathbb{Z}(n))$ has a basis consisting of common eigenvectors of \widehat{s}_n and \widehat{f}_n . Let $L_+(\mathbb{Z}(n))$ and $L_-(\mathbb{Z}(n))$ be the eigenspaces of the involution \widehat{f}_n^2 corresponding to the eigenvalues $+1$ and -1 , respectively. The action of \widehat{s}_n on $L(\mathbb{Z}(n))$ must respect the decomposition

$$L(\mathbb{Z}(n)) = L_+(\mathbb{Z}(n)) \oplus L_-(\mathbb{Z}(n)).$$

The sets $\{x^* + (-x)^* : x \in \mathbb{Z} \cap [0, n/2]\}$ and $\{x^* - (-x)^* : x \in \mathbb{Z} \cap (0, n/2)\}$ are bases for $L_+(\mathbb{Z}(n))$ and $L_-(\mathbb{Z}(n))$, respectively. With respect to these two bases, the actions of \widehat{s}_n on $L_+(\mathbb{Z}(n))$ and on $L_-(\mathbb{Z}(n))$ are both represented by tridiagonal matrices whose off-diagonal

entries are all non-zero. By Wilkinson [18, Section 5.38], these two actions of \widehat{s}_n are both separable. Therefore, up to scalar multiplication, $L_+(\mathbb{Z}(n)) \cup L_-(\mathbb{Z}(n))$ contains a unique basis of eigenvectors of \widehat{s}_n . The assertion follows because any eigenvector of \widehat{f}_n must belong either to $L_+(\mathbb{Z}(n))$ or else to $L_-(\mathbb{Z}(n))$. \square

By Remark 2.1, Theorem 2.2, and the fact that \widehat{s}_n is hermitian, there exists a unique basis S_n of $L(\mathbb{Z}(n))$ consisting of real unit eigenvectors σ such that either $\sigma(0) > 0$ or else $\sigma(0) = 0 < \sigma(1)$. Since the action of \widehat{s}_n on $L_+(\mathbb{Z}(n))$ is separable, we can enumerate the elements of $S_n \cap L_+(\mathbb{Z}(n))$ as $\sigma_{n,0}, \sigma_{n,2}, \sigma_{n,4}, \dots$ such that the corresponding eigenvalues are in strictly decreasing order. Similarly, we can enumerate the elements of $S_n \cap L_-(\mathbb{Z}(n))$ as $\sigma_{n,1}, \sigma_{n,3}, \sigma_{n,5}, \dots$ such that the corresponding eigenvalues are in strictly decreasing order. By considering the identity $n = |\mathbb{Z} \cap [0, n/2]| + |\mathbb{Z} \cap (0, n/2)|$, we see that $L_+(\mathbb{Z}(n))$ has dimension either $n/2 + 1$ or $(n + 1)/2$ (whichever is an integer), likewise $L_-(\mathbb{Z}(n))$ has dimension either $n/2 - 1$ or $(n - 1)/2$. Thus we have enumerated the elements $\sigma_{n,k}$ of S_n such that, if n is even, then the index k is an integer satisfying $0 \leq k \leq n - 2$ or $k = n$, while if n is odd, then k is an integer satisfying $0 \leq k \leq n - 1$. After Candan [7], we define, for each $t \in \mathbb{R}$, the **discrete fractional Fourier Transform** $\widehat{f}_n^{[t]}$ to be the operator on $L(\mathbb{Z}(n))$ such that

$$\widehat{f}_n^{[t]} \sigma_{n,k} := e^{2\pi i k t} \sigma_{n,k}$$

for each index k as above. Evidently, $\widehat{f}_n^{[t+t']} = \widehat{f}_n^{[t]} \widehat{f}_n^{[t']}$ for all $t, t' \in \mathbb{R}$. Note that $\widehat{f}_n^{[1]} = 1$. Furthermore, $\widehat{f}_n^{[1/4]}$ has order 4. Candan observed that, by McClellan–Parks [11], the multiplicities of $1, i, -1, -i$ as eigenvalues of $\widehat{f}_n^{[1/4]}$ coincide with the multiplicities of $1, i, -1, -i$, respectively, as eigenvalues of \widehat{f}_n . This, together with numerical evidence in Pei–Yeh [15] and Candan–Kutay–Özaktas [8] supports the following conjecture.

CONJECTURE 2.A: *Given an integer $n \geq 1$, then $\widehat{f}_n^{[1/4]} = \widehat{f}_n$.*

Theorems 2.6 and 2.7, below, tell us that $\widehat{f}_n^{[1/4]}$ and \widehat{f}_n both induce \widehat{f} .

Consider a vector $\phi \in \mathcal{S}(\mathbb{R})$. Given a vector $\phi_n \in L(\mathbb{Z}(n))$, we say that ϕ **restricts** to ϕ_n , writing $\phi_n = \text{res}_n(\phi)$, provided, for all $x \in \mathbb{Z}(n)$, we have

$$\phi(x\nu(n)^{-2}) = \nu(n)\phi_n(x).$$

Let \mathcal{N} be an infinite set of natural numbers such that, given $m, n \in \mathcal{N}$ with $n \leq m$, then n divides m , and the integer m/n is a square. The set

$$\mathbb{R}(\mathcal{N}) := \{x\nu(n)^{-2} : n \in \mathcal{N}, x \in \mathbb{Z}(n)\}$$

is dense in \mathbb{R} . Given an element $\xi \in \mathbb{R}(\mathcal{N})$, we define $\rho(\xi)$ to be the minimal element of \mathcal{N} such that $\xi\nu(\rho(\xi))^2 \in \mathbb{Z}(\rho(\xi))$. Thus, for each $n \in \mathcal{N}$ with $n \geq \rho(\xi)$, we can write

$$\xi = \xi(n)\nu(n)^{-2}$$

where $\xi(n) \in \mathbb{Z}(n)$. Let $L(\mathbb{Z}(\mathcal{N}))$ denote the space of sequences of the form $(\psi_n : n \in \mathcal{N})$ where each $\psi_n \in L(\mathbb{Z}(n))$. Since $\mathbb{R}(\mathcal{N})$ is dense in \mathbb{R} , each vector $\psi \in \mathcal{S}(\mathbb{R})$ is uniquely determined by the element $(\text{res}_n(\psi) : n \in \mathcal{N})$ of $L(\mathbb{Z}(\mathcal{N}))$. Note that any infinite subset \mathcal{N}' of \mathcal{N} satisfies the hypothesis we imposed upon \mathcal{N} , and furthermore, $\mathbb{R}(\mathcal{N}') = \mathbb{R}(\mathcal{N})$.

The condition on \mathcal{N} that each integer m/n be a square is not essential, and can be dispensed with, but at the price of introducing some irritating complications that would obscure our

arguments. We do not know whether our results would still hold if \mathcal{N} were replaced with any infinite subset of \mathbb{N} . Anyway, the hypothesis we have imposed upon \mathcal{N} covers all the applications we have in mind. As an example, we can take \mathcal{N} to consist of all the positive even powers of some given integer $d \geq 2$, or we can take \mathcal{N} to consist of all the positive odd powers of d . The case where d is prime pertains to the references to discrete quantum mechanics cited in Section 1.

Consider a vector $\psi \in L^2(\mathbb{R})$, and a sequence $(\psi_n : n \in \mathcal{N}) \in L(\mathbb{Z}(\mathcal{N}))$. We say that $(\psi_n : n \in \mathcal{N})$ **induces** ψ , writing $\psi = \text{ind}_{n \in \mathcal{N}}(\psi_n)$, provided, for all $\phi \in \mathcal{S}(\mathbb{R})$, we have

$$\langle \phi | \psi \rangle = \lim_{n \in \mathcal{N}} \langle \text{res}_n(\phi) | \psi_n \rangle .$$

By the Riesz Representation Theorem, each element of $L(\mathbb{Z}(\mathcal{N}))$ induces at most one vector in $L^2(\mathbb{R})$.

LEMMA 2.3: *Given uniformly continuous vectors $\phi, \psi \in L^2(\mathbb{R})$, then*

$$\langle \phi | \psi \rangle = \lim_{n \in \mathcal{N}} \langle \text{res}_n(\phi) | \text{res}_n(\psi) \rangle .$$

Proof: Let $\phi_n := \text{res}_n(\phi)$ and $\psi_n := \text{res}_n(\psi)$. Since ϕ and ψ are Riemann-integrable,

$$\int_{-q}^q \phi(r)^* \psi(r) dr = \lim_{n \in \mathcal{N}} \sum_{x \in \mathbb{Z}(\mathcal{N}) : |x| \leq q\nu(n)^2} \phi_n(x)^* \psi_n(x)$$

for all real $q > 0$. Via the Cauchy–Schwartz inequality, we easily obtain

$$\left(\lim_{n \in \mathcal{N}} \sum_{x \in \mathbb{Z}(n) : |x| > q\nu(n)^2} \phi_n(x)^* \psi_n(x) \right)^2 \leq \int_{r: |r| \geq q} |\phi(r)|^2 dr \cdot \int_{r: |r| \geq q} |\psi(r)|^2 dr .$$

The right-hand expression tends to zero as q tends to infinity. \square

LEMMA 2.4: *Given a uniformly continuous vector $\psi \in L^2(\mathbb{R})$, then $\psi = \text{ind}_{n \in \mathcal{N}}(\text{res}_n(\psi))$.*

Proof: This is immediate from Lemma 2.3. \square

The four Theorems stated below will be proved in Section 3.

THEOREM 2.5: *For each $j \in \mathbb{N}$, we have $h_j = \text{ind}_{n \in \mathcal{N}}(\sigma_{n,j})$.*

THEOREM 2.6: *Given a vector $\psi \in L^2(\mathbb{R})$, then there exists an element $(\psi_n : n \in \mathcal{N}) \in L(\mathbb{Z}(\mathcal{N}))$ such that $\psi = \text{ind}_{n \in \mathcal{N}}(\psi(n))$.*

For $n \in \mathcal{N}$, let $\text{End}(L(\mathbb{Z}(n)))$ denote the ring of operators on $L(\mathbb{Z}(n))$. The ring

$$\text{End}(L(\mathbb{Z}(\mathcal{N}))) := \prod_{n \in \mathcal{N}} \text{End}(L(\mathbb{Z}(n)))$$

consists of the sequences $(\hat{a}_n : n \in \mathcal{N})$ such that each $\hat{a}_n \in \text{End}(L(\mathbb{Z}(n)))$. We say that $(\hat{a}_n : n \in \mathcal{N})$ **induces** \hat{a} , writing $\hat{a} = \text{ind}_{n \in \mathcal{N}}(\hat{a}_n)$, provided, given any element $(\psi_n : n \in \mathcal{N}) \in L(\mathbb{Z}(\mathcal{N}))$ inducing a vector $\psi \in L^2(\mathbb{R})$, then $(\hat{a}_n \psi_n : n \in \mathcal{N})$ induces $\hat{a}\psi$. The Riesz Representation Theorem, together with Theorem 2.6, ensures that each element of $\text{End}(L(\mathbb{Z}(\mathcal{N})))$ induces at most one operator on $L^2(\mathbb{R})$.

The condition that $(\widehat{a}_n : n \in \mathcal{N})$ induces \widehat{a} seems to be rather strong. Suppose, for instance, that each a_n is given by

$$\widehat{a}_n x^* = \nu(n)^2((x-1)^* - 2x^* + (x+1)^*).$$

In numerical calculation, \widehat{a}_n is often used as a discrete approximation to the second differential operator. Of course, differential operators do not have domain $L^2(\mathbb{R})$, but that is not an essential problem, because we could easily extend the definition of induction so as to be applicable to operators on a domain dense in $L^2(\mathbb{R})$. Our point is that, putting $\psi_n(x) = (-1)^x n^\gamma$ with $\gamma \in \mathbb{R}$, then $\widehat{a}_n \psi_n = -4\nu(n)^2 \psi_n$, and there clearly exist γ for which $(\psi_n : n \in \mathcal{N})$ induces the zero-vector while $(\widehat{a}_n \psi_n : n \in \mathcal{N})$ does not induce any vector. As another example, the operator \widehat{h}_n on $L(\mathbb{Z}(n))$ given by

$$\widehat{h}_n x^* = \nu(n)^2(-(x-1)^* + (4 - 2\cos(2\pi x/n))x^* - (x+1)^*)$$

may seem sensible as a discrete analogue of the operator \widehat{h} on $\mathcal{S}(\mathbb{R})$, but there does not exist a vector $\phi \in L^2(\mathbb{R})$ such that $(\widehat{h}_n \phi_n : n \in \mathcal{N})$ induces a vector for all $(\phi_n : n \in \mathcal{N}) \in L(\mathbb{Z}(\mathcal{N}))$ inducing ϕ .

THEOREM 2.7: *We have $\widehat{f} = \text{ind}_{n \in \mathcal{N}}(\widehat{f}_n)$.*

The essence of Theorem 2.7 has been known since the 1930's. We have been unable to derive it from any results we have found in the literature, but we suspect that it would yield to general principles of a Hilbert space approach to spectral methods as in, for instance, Picard [16, Section 2.2].

THEOREM 2.8: *For all $t \in \mathbb{R}$, we have $\widehat{f}^{[t]} = \text{ind}_{n \in \mathcal{N}}(\widehat{f}_n^{[t]})$.*

3: Proofs

For each $j \in \mathbb{N}$ and $n/in\mathcal{N}$, let $h_{n,j} := \text{res}_n(h_j)$ and $\lambda_{n,j} := 4 - 2\pi(2j+1)/n$.

LEMMA 3.1: *Given $j \in \mathbb{N}$, and $\xi \in \mathbb{R}(\mathcal{N})$ and $n \in \mathcal{N}$ with $n \geq \rho(\xi)$, then*

$$|((\widehat{s}_n - \lambda_{n,j})h_{n,j})(\xi(n))| = O(n^{-9/4}).$$

Proof: Let $x := \xi(n)$. Taylor's Theorem gives

$$h_j(\xi + \delta) = \sum_{\alpha=0}^{\infty} \delta^\alpha h_j^{(\alpha)}(\xi)/\alpha!.$$

Putting $\delta = \pm\nu(n)^{-2}$, and using the fact that each function $r \mapsto |h_j^{(\alpha)}(r)|$ is bounded above, we have

$$\begin{aligned} & h_{n,j}(x-1) - 2h_{n,j}(x) + h_{n,j}(x+1) \\ &= \nu(n)^{-5} h_j''(\xi) + \sum_{\alpha=4}^{\infty} (1 + (-1)^\alpha) \nu(n)^{-2\alpha-1} h_j^{(\alpha)}(\xi)/\alpha! = \nu(n)^{-5} h_j''(\xi) + O(\nu(n)^{-9}). \end{aligned}$$

Similar manipulations give $(-2 + 2\cos 2\pi x/n)h_{n,j}(x) = -\xi^2 \nu(n)^{-5} h_j(\xi) + O(\nu(n)^{-9})$, whence

$$((\widehat{s}_n - \lambda_{n,j})h_{n,j})(x) + O(n^{-9/4}) = \nu(n)^{-5}(h_j''(\xi) + (-\xi^2 + 2j + 1)h_j(\xi)) = 0. \quad \square$$

Given any vector v in an inner product space, we write the modulus of v as $\|v\|$. In particular, for $\psi \in L^2(\mathbb{R})$, we write $\|\psi\| := \|\psi\|_2$.

LEMMA 3.2: *Given $j \in \mathbb{N}$ and $n \in \mathcal{N}$, then $\|(\widehat{s}_n - \lambda_{n,j})h_{n,j}\| = o(n^{-3/2})$.*

Proof: Let $k_n := (\widehat{s}_n \lambda_{n,j})h_{n,j}$. By Lemma 3.1,

$$\sum_{x \in \mathbb{Z}(n); |x| \leq \nu(n)^2 \log n} |k_n(x)|^2 = O(n^{-4} \log n).$$

Choose $B \in \mathbb{R}$ such that $|h_j(r)| < Br^j e^{-r^2/2}$ for all $r \in \mathbb{R}$. For $x \in \mathbb{Z}(n)$ and $\xi = x\nu(n)^{-2}$, we have $\nu(n)|k_n(x)| \leq 8B\xi^j e^{-\xi^2/2}$ because $|\lambda_{n,j}| \leq 4$ and \widehat{s}_n is a sum of four unitary operators. So

$$\sum_{x \in \mathbb{Z}(n); |x| > \nu(n)^2 \log n} |k_n(x)|^2 \leq 64B^2 \int_{r: |r|+1 \geq \log n} r^{2j} e^{-r^2} dr.$$

The right-hand expression diminishes faster than $n^{-4} \log n$, so $\|k_n\| = o(n^{-2} \log n)$. \square

LEMMA 3.3: *Given $j \in \mathbb{N}$, and sufficiently large $n \in \mathcal{N}$, then \widehat{s}_n has an eigenvalue in the open interval $(4 - 4\pi j/n - 3\pi/n, 4 - 4\pi j/n - \pi/n)$.*

Proof: Suppose, for a contradiction, that there exists an infinite subset \mathcal{N}' of \mathcal{N} such that, for each $n \in \mathcal{N}'$, there is no eigenvalue of \widehat{s}_n in the specified interval. Since the function h_j has precisely j zeroes, we cannot have $\|h_{n,j}\| = 0$ unless $n \leq j$. So we may define $g_{n,j} := h_{n,j} / \|h_{n,j}\|$ for all $n \in \mathcal{N}'$ with $n > j$. Lemma 2.3 gives $\lim_{n \in \mathcal{N}'} \|h_{n,j}\| = \|h_j\| = 1$, and then Lemma 3.2 tells us that

$$\|(\widehat{s}_n - (4 - 4\pi j/n - 2\pi/n))g_{n,j}\| = o(n^{-3/2}).$$

For each $n \in \mathcal{N}'$ with $n > j$, the hypothesis on \mathcal{N}' allows us to write $g_{n,j} = a_n + b_n$, where a_n is a linear combination of eigenvectors of \widehat{s}_n with associated eigenvalues at least $4 - 4\pi j/n - \pi/n$, and b_n is a linear combination of eigenvectors of \widehat{s}_n with associated eigenvalues at most $4 - 4\pi j/n - 3\pi/n$. Using the fact that a_n and b_n are orthogonal, we obtain the contradiction

$$\|(\widehat{s}_n - (4 - 4\pi j/n - 2\pi/n))(a_n + b_n)\| \geq \pi/n(\|a_n\|^2 + \|b_n\|^2) = \pi/n. \quad \square$$

The next five results concern the following scenario: let $F \in \mathbb{R}$, and for each $n \in \mathcal{N}$, let ω_n be a real unit eigenvector of both \widehat{s}_n and \widehat{f}_n such that the eigenvalue of \widehat{s}_n associated with ω_n has the form $4 - 2\pi E_n/n$ where $E_n \leq F$. Thus ω_n satisfies Harper's Equation

$$\omega_n(x-1) + \omega_n(x+1) = 2(1 - \pi E_n/n - \cos 2\pi x/n)\omega_n(x)$$

for $x \in \mathbb{Z}$. Since $\widehat{s}_n = \widehat{u}_n + \widehat{u}_n^\dagger + \widehat{v}_n + \widehat{v}_n^\dagger$ as a sum of four unitary operators which do not mutually commute, we have $E_n > 0$, and in particular, $F > 0$.

LEMMA 3.4: *Let $n \in \mathcal{N}$, and let $x \in \mathbb{Z}(n)$ such that $\cos 2\pi x/n \leq 1 - \pi E_n/n$. If $x > 0$, then $|\omega_n(x)| \leq |\omega_n(x-1)|$. If $x < 0$, then $|\omega_n(x)| \leq |\omega_n(x+1)|$.*

Proof: Since ω_n is an eigenvector of \widehat{f}_n^2 , we have $|\omega_n(z)| = |\omega_n(-z)|$ for all $z \in \mathbb{Z}$. So, if the assertion fails, then there must exist some $x \in \mathbb{Z}$ such that $\cos 2\pi x/n \leq 1 - \pi E_n/n$ and $|\omega_n(x-1)| < |\omega_n(x)| > |\omega_n(x+1)|$. But Harper's Equation forces $|\omega_n(x-1) + \omega_n(x+1)| \geq 2|\omega_n(x)|$. \square

LEMMA 3.5: *Given sufficiently large $n \in \mathcal{N}$, and $x \in \mathbb{Z}(n)$, then $\nu(n)|\omega_n(x)| \leq (2F)^{1/4}$.*

Proof: For each $n \in \mathcal{N}$, let M_n be the maximum value of $|\omega_n(x)|$ with $x \in \mathbb{Z}$. Let us fix an element $n \in \mathcal{N}$, and assume that n is sufficiently large for our purposes. Write $M := M_n$, and choose $y \in \mathbb{Z}(n)$ such that $\omega_n(y) = M$. Since $|\omega_n(y)| = |\omega_n(-y)|$, we may insist that $y < 0$. We assume that $\omega_n(y) > 0$; the argument when $\omega_n(y) < 0$ is similar. Since n is sufficiently large, Lemma 3.4 and the condition $y < 0$ ensure that

$$1 \geq \cos 2\pi(y+x)/n > 1 - \pi E_n/n \geq 1 - \pi F/n$$

for each integer z with $0 \leq z \leq \sqrt{n/\pi F}$. For each such z , let $\Delta(z) := \omega_n(y+z) - \omega_n(y+z-1)$. We have $|\Delta(z+1) - \Delta(z)| \leq 2\pi FM/n$, hence $|\Delta(z)| \leq 2\pi FMz/n$ and

$$\omega_n(y+z) \geq M - \sum_{x=0}^z 2\pi FMx/n = M(1 - \pi Fz(z+1)/n) > M(1 - \pi F(z+1)^2/n).$$

Letting m be the largest integer such that $m \leq \sqrt{n/\pi F}$, then

$$\begin{aligned} 1 &= \|\omega_n\|^2 \geq \sum_{z=0}^{m-1} \omega_n(y+z)^2 > M^2 \sum_{z=1}^m (1 - \pi Fz^2/n)^2 \\ &= M^2 \left(m - \frac{2\pi F}{n} \left(\frac{m^3}{3} + \frac{m^2}{2} + \frac{m}{6} \right) + (\pi F/n)^2 \left(\frac{m^5}{5} + \frac{m^4}{2} + \frac{m^3}{3} - \frac{m}{30} \right) \right). \end{aligned}$$

Now letting the element $n \in \mathcal{N}$ vary, we have

$$1 > M_n^2 (8\sqrt{n/\pi F}/15 + o(\sqrt{n})) > M_n^2 (\nu(n)^2/\sqrt{2F} + o(\nu(n)^2)). \quad \square$$

We now construct a function $\omega_{\mathcal{N}} : \mathbb{R}(\mathcal{N}) \rightarrow \mathbb{R}$. Let n_0 be the smallest element of \mathcal{N} . Given an element $\zeta \in \mathbb{R}(\mathcal{N})$ with $\rho(\zeta) \leq n_0$, then Lemma 3.5 guarantees that the set $\{\nu(n)\omega_n(\zeta(n)) : n \in \mathcal{N}\}$ has a cluster point $\omega_{\mathcal{N}}(\zeta)$. Since there are only finitely many $\zeta \in \mathbb{R}(\mathcal{N})$ with $\rho(\zeta) \leq n_0$, there exists an infinite subset \mathcal{N}_1 of \mathcal{N} such that $n_0 \notin \mathcal{N}_1$ and

$$\omega_{\mathcal{N}}(\zeta) = \lim_{n \in \mathcal{N}_1} (\nu(n)\omega_n(\zeta(n)))$$

for all $\zeta \in \mathbb{R}(\mathcal{N})$ with $\rho(\zeta) \leq n_0$.

Now let $t \geq 1$, and assume, inductively, that we have defined \mathcal{N}_s for all integers s with $1 \leq s \leq t$. For each s , let n_s be the smallest element of \mathcal{N}_s . We also assume that we have defined $\omega_{\mathcal{N}}(\zeta)$ for all $\zeta \in \mathbb{R}(\mathcal{N})$ with $\rho(\zeta) \leq n_{t-1}$. As before, there exists an infinite subset \mathcal{N}_{t+1} of \mathcal{N}_t such that $n_t \notin \mathcal{N}_{t+1}$, and the real number

$$\omega_{\mathcal{N}}(\zeta) = \lim_{n \in \mathcal{N}_{t+1}} (\nu(n)\omega_n(\zeta(n)))$$

is a cluster point of $\{\nu(n)\omega_n(\zeta(n)) : n \in \mathcal{N}_t\}$ for all $\zeta \in \mathbb{R}(\mathcal{N})$ with $\rho(\zeta) \leq n_t$. Letting $\mathcal{N}_* := \{n_t : t \in \mathcal{N}\}$, then

$$\omega_{\mathcal{N}}(\zeta) = \lim_{n \in \mathcal{N}_*} (\nu(n)\omega_n(\zeta(n)))$$

for all $\zeta \in \mathbb{R}(\mathcal{N})$.

LEMMA 3.6: *Given $\eta \in \mathbb{R}(\mathcal{N})$ with $\eta^2 \geq 4F$, then $\omega_{\mathcal{N}}(\eta)^2 \leq 4F/|\eta|^3$.*

Proof: Given real $r > 0$, and $n \in \mathcal{N}$, let $\Omega_n(r) := \sum_{x \in Z} \omega_n(x)^2$ and $\Omega'_n(r) := \sum_{x \in Z'} \omega_n(x)^2$, where $Z := \{x \in \mathbb{Z}(n) : |x| \leq r\nu(n)^2\}$ and $Z' := \{x \in \mathbb{Z}(n) : |x| > r\nu(n)^2\}$. Then

$$\Omega_n(r) + \Omega'_n(r) = \|\omega_n\|^2 = 1.$$

Now $\widehat{s}_n = \widehat{u}_n + \widehat{u}_n^\dagger + \widehat{v}_n + \widehat{v}_n^\dagger$ as a sum of four unitary operators, so

$$4 - 2\pi F/n \leq \langle \omega_n | \widehat{s}_n | \omega_n \rangle = \langle \omega_n | \widehat{u}_n + \widehat{u}_n^\dagger | \omega_n \rangle + \langle \omega_n | \widehat{v}_n + \widehat{v}_n^\dagger | \omega_n \rangle$$

and both of the terms of the sum have absolute value at most 2. Therefore

$$\begin{aligned} 1 - \pi F/n &\leq \langle \omega_n | \widehat{u}_n + \widehat{u}_n^\dagger | \omega_n \rangle / 2 = \sum_{x \in \mathbb{Z}(n)} \omega_n(x)^2 \cos 2\pi x/n \\ &\leq \Omega_n(r) + \Omega'_n(r) \cos 2\pi r\nu(n)^2/n = 1 - \Omega'_n(r) \pi r^2/n + O(n^{-2}). \end{aligned}$$

We deduce that $\Omega'_n(r) < F/r^2 + O(1/n)$. We may assume that $\eta > 0$. Let $n \in \mathcal{N}$ such that $n \geq \rho(\eta)$. Writing $y := \eta(n)$, then

$$\cos \pi y/n + O(n^{-2}) < 1 - \pi \eta^2/4n \leq 1 - \pi E_n/n.$$

By Lemma 3.4, $|\omega_n(x)|$ is monotonically decreasing in the range $x \in \mathbb{Z} \cap [y/2, y]$. Recalling that $\omega_n(x)^2 = \omega_n(-x)^2$, and summing $\omega_n(x)^2$ over the integers $x \in [-y, -y/2] \cup [y/2, y]$, we obtain $\omega_n(y)^2(y + O(1)) \leq \Omega'_n(\eta/2)$. Lemma 3.5 implies that $\omega_n(y)^2 = O(n^{-1/2})$, so

$$\omega_n(y)^2 \eta \nu(n)^2 + O(n^{-1/2}) < 4F/\eta^2.$$

The assertion now follows from the construction of $\omega_{\mathcal{N}}$. \square

We allow $\mathbb{R}(\mathcal{N})$ to inherit the subspace topology from \mathbb{R} .

LEMMA 3.7: *The function $\omega_{\mathcal{N}} : \mathbb{R}(\mathcal{N}) \rightarrow \mathbb{R}$ is continuous.*

Proof: Let $\xi, \eta \in \mathbb{R}(\mathcal{N})$ with $\xi \geq \eta$. We shall calculate an upper bound for $\omega_{\mathcal{N}}(\xi) - \omega_{\mathcal{N}}(\eta)$. Let $n \in \mathcal{N}$, and assume that n is large; in particular, $\rho(\xi) \leq n \leq \rho(\eta)$. Let $x := \xi(n)$ and $y := \eta(n)$. The eigenvalue i_n of \widehat{f}_n associated with ω_n is a 4-th root of unity, and

$$\begin{aligned} i_n(\omega_n(x) - \omega_n(y)) &= (\widehat{f}_n \omega_n)(x) - (\widehat{f}_n \omega_n)(y) = \frac{1}{\sqrt{n}} \sum_{z \in \mathbb{Z}(n)} \omega_n(z) (e^{2\pi i x z/n} - e^{2\pi i y z/n}) \\ &= \frac{-2i}{\sqrt{n}} \sum_{z \in \mathbb{Z}(n)} \omega_n(z) e^{\pi i (x+y)z/n} \sin \pi (x-y)z/n. \end{aligned}$$

Therefore

$$\nu(n)^2 |\omega_n(x) - \omega_n(y)| \leq \sqrt{\frac{2}{\pi}} \sum_{z \in \mathbb{Z}(n)} |\omega_n(z) \sin \pi (x-y)z/n|.$$

Let $\tau \in \mathbb{R}(\mathcal{N})$ such that $\tau \geq 2\sqrt{F}$. Insisting that $n \geq \rho(\tau)$, let $t := \tau(n)$. Let $Z := \{x \in \mathbb{Z} : |x| \leq t\}$, and $Z' := \mathbb{Z}(n) - Z$. Bearing in mind that n is large, the proof of Lemma 3.6 shows that

$$|\omega_n(z)| < \sqrt{5F/\zeta^3}/\nu(n) = \nu(n)^2 \sqrt{5F} z^{-3/2}$$

where $z \in Z'$ and $\zeta := z\nu(n)^{-2}$. Therefore

$$\begin{aligned} \sum_{z \in Z'} |\omega_n(z) \sin \pi(x-y)z/n| &< 2\nu(n)^2 \sqrt{5F} \sum_{z=t+1}^{\infty} z^{-3/2} \\ &\leq 6\nu(n)^2 \sqrt{F} \int_t^{\infty} r^{-3/2} dr = 12\nu(n)^2 \sqrt{F/t} = 12\nu(n) \sqrt{F/\tau}. \end{aligned}$$

On the other hand, via Lemma 3.5,

$$\begin{aligned} \sum_{z \in Z} |\omega_n(z) \sin \pi(x-y)z/n| &\leq \frac{(2F)^{1/4}}{\nu(n)} \sum_{z=-t}^t |\pi(x-y)z/n + O(n^{-3/2})| \\ &\leq (2F)^{1/4} (\pi(x-y)t(t+1)/n + O(n^{-1/2}))/\nu(n) < F^{1/4} ((x-y)t^2/\nu(n) + O(n^{-3/2})) \\ &= F^{-1/4} \nu(n) \tau^2 (\zeta - \eta) + O(n^{-3/2}). \end{aligned}$$

Therefore, $\nu(n)|\omega_n(x) - \omega_n(y)| \leq \sqrt{2/\pi} (F^{1/4} \tau^2 (\zeta - \eta) + 12\sqrt{F/\tau}) + O(1/n)$, hence

$$|\omega_{\mathcal{N}}(\zeta) - \omega_n(\eta)| < F^{1/4} \tau^2 (\zeta - \eta) + 12\sqrt{F/\tau}. \quad \square$$

LEMMA 3.8: *The function $\omega_{\mathcal{N}} : \mathbb{R}(\mathcal{N}) \rightarrow \mathbb{R}$ extends uniquely to a real-valued continuous function $\omega : \mathbb{R} \rightarrow \mathbb{C}$. Furthermore, ω is a uniformly continuous function in $L^2(\mathbb{R})$, and $\|\omega\| = 1$.*

Proof: The first part is immediate from Lemma 3.7. By Lemmas 3.5 and 3.6, the function $r \mapsto |\omega(r)|$ is bounded by both $(2F)^{1/4}$ and $O(|r|^{-3/2})$. So ω is uniformly continuous.

To prove the last part, we may assume that $\mathcal{N} = \mathcal{N}^*$. Let $\tau \in \mathbb{R}(\mathcal{N})$ with $\tau \geq \sqrt{4F}$. Lemma 3.6 gives $\omega(r)^2 \leq 4F/|r|^3$ whenever $|r| \geq \tau$. In particular,

$$\lim_{\tau \rightarrow \infty} \int_{|r| > \tau} \omega(r)^2 dr = 0.$$

The proof of Lemma 3.6 shows that

$$\lim_{\tau \rightarrow \infty} \lim_{n \in \mathcal{N}} \sum_{x \in \mathbb{Z}(n) : |x| > \tau \nu(n)^2} \omega_n(x)^2 = 0.$$

Let us now fix τ . Since each $\|\omega_n\| = 1$, it suffices to show that

$$\int_{-\tau}^{\tau} \omega(r)^2 dr = \lim_{n \in \mathcal{N}} \sum_{x \in Z} \omega_n(x)^2$$

where $Z := \{x \in \mathbb{Z}(n) : |x| \leq \tau \nu(n)^2\}$. Choose any real $\epsilon > 0$. The proof of Lemma 3.7 shows that, given sufficiently large $n \in \mathcal{N}$, then for all $\xi, \xi' \in \mathbb{R}(\mathcal{N})$ with $|\xi - \xi'| < \delta$, we have

$$|\nu(n)^2 (\omega_n(\xi(n))^2 - \omega(\xi'(n))^2)| < \epsilon/2.$$

For each $\zeta \in \mathbb{R}(\mathcal{N}) \cap [-\tau, \tau]$, we have

$$|\nu(n)^2 \omega_n(\zeta(n))^2 - \omega(\zeta)^2| < \epsilon/2$$

for sufficiently large n . So, for each such ζ , there exists some neighbourhood $U(\zeta)$ of ζ in $[-\tau, \tau]$ such that, for all $r \in U(\zeta)$, we have

$$|\omega(r)^2 - \omega(\zeta)^2| < \epsilon$$

and furthermore, given sufficiently large n , then for all $\xi \in \mathbb{R}(\mathcal{N}) \cap U(\zeta)$ with $\rho(\xi) \leq n$, we have

$$|\nu(n)^2 \omega_n(\xi(n))^2 - \omega(\zeta)^2| < \epsilon.$$

Let $\{U(\zeta_\alpha) : \alpha \in A(\epsilon)\}$ be a finite subcover of $[-\tau, \tau]$, and write $[-\tau, \tau] = \bigcup_{\alpha \in A(\epsilon)} V(\epsilon, \alpha)$, as a disjoint union, where each $V(\epsilon, \alpha)$ is a connected subset of $U(\zeta_\alpha)$. We have a Lebesgue integral

$$\int_{-\tau}^{\tau} \omega(r)^2 dr = \lim_{\epsilon \rightarrow 0} \sum_{\alpha \in A(\epsilon)} \mu(V(\epsilon, \alpha)) \omega(\zeta_\alpha)^2.$$

where μ denotes the Borel measure. Since there are $O(\sqrt{n})$ elements x of $\mathbb{Z}(n)$ satisfying $|x| \leq \tau \nu(n)^2$, we have

$$\sum_{\alpha \in A(\epsilon)} \mu(V(\epsilon, \alpha)) \omega(\zeta_\alpha)^2 + \epsilon O(1) = \sum_{\alpha \in A(\epsilon)} \left(\sum_{x \in \mathbb{Z}(n): x \nu(n)^{-2} \in V(\epsilon, \alpha)} \omega_n(x)^2 \right) = \sum_{x \in \mathbb{Z}} \omega_n(x)^2$$

for sufficiently large n . \square

LEMMA 3.9: *Given $j \in \mathbb{N}$, and sufficiently large $n \in \mathcal{N}$, then \hat{s}_n has a unique eigenvalue in the closed interval $[4 - 4\pi(j+1)/n, 4 - 4\pi j/n]$.*

Proof: Thanks to Lemma 3.3, it suffices to derive a contradiction from the assumption that, for infinitely many $n \in \mathcal{N}$, there exist two orthogonal real unit eigenvectors ω'_n and ω''_n of \hat{s}_n whose associated eigenvalues lie in the specified interval. We replace \mathcal{N} with the set of such n . For each $n \in \mathcal{N}$, we put either $\omega_n = \omega'_n$ or else $\omega_n = \omega''_n$, insisting that $\langle \omega_n | h_{n,j} \rangle \leq 3/4$ for sufficiently large n . By Lemma 3.2, $\lim_{n \in \mathcal{N}} \langle \omega_n | h_{n,j'} \rangle^2 = 0$ for all $j' \in \mathbb{N}$ with $j' \neq j$. Since $\{h_j : j \in \mathbb{N}\}$ is an orthonormal basis for $L^2(\mathbb{R})$, we have $\sum_{k \in \mathbb{N}} \langle \omega | h_k \rangle^2 = 1$. The desired contradiction will be achieved when we have shown that

$$\langle \omega | h_k \rangle = \lim_{n \in \mathcal{N}} \langle \omega_n | h_{n,k} \rangle.$$

The argument is similar to the proof of Lemma 3.8; we shall outline the steps.

Using Lemma 3.6, and the fact that h_j is rapidly decreasing, it is easy to reduce to the task of proving that

$$\int_{-\tau}^{\tau} \omega(r) h_k(r) dr = \lim_{n \in \mathcal{N}} \sum_{x \in \mathbb{Z}} \omega_n(x) h_{n,j}(x)$$

where τ and Z are as in the proof of Lemma 3.8. For each $\zeta \in \mathbb{R}(\mathcal{N}) \cap [-\tau, \tau]$, we choose a neighbourhood $U'(\zeta)$ of ζ such that, for all $r \in U'(\zeta)$, we have

$$|\omega(r) h_k(r) - \omega(\zeta) h_k(\zeta)| < \epsilon$$

and furthermore, given sufficiently large $n \in \mathcal{N}$, then for all $\xi \in \mathbb{R}(\mathcal{N}) \cap U(\zeta)$ with $\rho(\xi) \leq n$, we have

$$|\nu(n)^2 \omega_n(\xi(n)) h_{n,k}(\xi(n)) - \omega(\zeta) h_k(\zeta)| < \epsilon.$$

Choosing a finite subcover $\{U'(\zeta'_\alpha) : \alpha \in A'(\epsilon)\}$, and writing $[-\tau, \tau] = \bigcup_{\alpha \in A'(\epsilon)} V'(\epsilon, \alpha)$ just as before, we have

$$\sum_{\alpha \in A'(\epsilon)} \mu(V'(\epsilon, \alpha)) \omega(\zeta'_\alpha) h_k(\zeta'_\alpha) + \epsilon O(1) = \sum_{x \in \mathbb{Z}} \omega_n(x) h_{n,k}(x).$$

Again, the assertion follows by letting $n \rightarrow \infty$ and then letting $\epsilon \rightarrow 0$. \square

Proof of Theorem 2.5: For each $n \in \mathcal{N}$, let us enumerate the elements $\sigma'_{n,0}, \sigma'_{n,1}, \dots, \sigma'_{n,n-1}$ of the basis S_n such that the associated eigenvalues $\lambda'_{n,0}, \lambda'_{n,1}, \dots, \lambda'_{n,n-1}$ are in decreasing order (perhaps not strictly). Lemmas 3.3 and 3.9 imply that, given any $j \in \mathbb{N}$, then for sufficiently large $n \in \mathcal{N}$, we have

$$4 - 4\pi j/n - 3\pi/n < \lambda'_{n,j} < 4 - 4\pi j/n - \pi/n.$$

Lemma 3.9 also implies that, for such given j and chosen n , we may write $h_{n,j} = \alpha_{n,j} + c_{n,j} \sigma'_{n,j} + \beta_{n,j}$ where $c_{n,j} \in \mathbb{C}$, and $\alpha_{n,j}$ is a linear combination of eigenvectors of \widehat{s}_n with associated eigenvalues at least $4 - 4\pi j/n$, and $\beta_{n,j}$ is a linear combination of eigenvectors of \widehat{s}_n with associated eigenvalues at most $4 - 4\pi(j+1)/n$. We saw in the proof of Lemma 3.3 that $\|h_{n,j}\| \rightarrow 1$ as n increases, so Lemma 3.2 gives $\lim_{n \in \mathcal{N}} \|\alpha_{n,j}\| = \lim_{n \in \mathcal{N}} \|\beta_{n,j}\| = 0$, and $\lim_{n \in \mathcal{N}} |c_{n,j}| = 1$. But $h_{n,j}$ and $\sigma'_{n,j}$ both have real coordinates with respect to the bases $\{x^* : x \in \mathbb{Z}(n)\}$ of $L(\mathbb{Z}(n))$, furthermore, if j is even, then $h_{n,j}(0) > 0$, while if j is odd, then $h_{n,j}(0) = 0 < h_{n,j}(1)$. Therefore $\lim_{n \in \mathcal{N}} c_{n,j} = 1$. For fixed $j \in \mathbb{N}$ and sufficiently large $n \in \mathcal{N}$, we have shown that $\sigma'_{n,j} = \sigma_{n,j}$. It follows that, when n is sufficiently large, $\sigma_{n,j} = (h_{n,j} - \alpha_{n,j} - \beta_{n,j})/c_{n,j}$. Given any uniformly continuous $\phi \in L^2(\mathbb{R})$, Lemma 2.3 now gives

$$\lim_{n \in \mathcal{N}} \langle \text{res}_n(\phi) | \sigma_{n,j} \rangle = \lim_{n \in \mathcal{N}} \langle \text{res}_n(\phi) | h_{n,j} \rangle = \langle \phi | h_j \rangle. \quad \square$$

Proof of Theorem 2.6: The proof of Theorem 2.5 shows that

$$\lim_{n \in \mathcal{N}} \langle h_{n,j} \sigma_{n,k} \rangle = \delta_{j,k}$$

for all $j, k \in \mathbb{N}$. Write $\psi = \sum_{j \in \mathbb{N}} c_j h_j$ with each $c_j \in \mathbb{C}$. For each $n \in \mathcal{N}$, let $\psi_n := \sum_{j=0}^{n-2} c_j \sigma_{n,j}$. Let $\mathcal{F}(\mathbb{R})$ denote the subspace of $\mathcal{S}(\mathbb{R})$ consisting of the linear combinations $\theta = \sum_{j \in \mathbb{N}} d_j h_j$ with only finitely many of the d_j non-zero. For each such θ , we have

$$\langle \theta | \psi \rangle = \sum_{j \in \mathbb{N}} d_j c_j = \lim_{n \in \mathcal{N}} \langle \text{res}_n(\theta) | \psi_n \rangle.$$

The assertion follows because $\mathcal{F}(\mathbb{R})$ is dense in the space of uniformly continuous functions in $L^2(\mathbb{R})$. \square

Proof of Theorem 2.7: This is similar to the proof, below, of Theorem 2.8. \square

Proof of Theorem 2.8: Let $\phi, \psi \in L^2(\mathbb{R})$ such that ϕ is uniformly continuous. Let $(\psi_n : n \in \mathcal{N}) \in L(\mathbb{Z}(\mathcal{N}))$ such that $\psi = \text{ind}_{n \in \mathcal{N}}(\psi_n)$. Let $t \in \mathbb{R}$, and write $\widehat{g} := \widehat{f}^{[t]}$. For each $n \in \mathcal{N}$, write $\widehat{g}_n := \widehat{f}_n^{[t]}$. We are to prove that

$$\langle \phi | \widehat{g}\phi \rangle = \lim_{n \in \mathcal{N}} \langle \text{res}_n(\phi) | \widehat{g}_n \psi_n \rangle.$$

Since $\mathcal{F}(\mathbb{R})$ is dense in $L^2(\mathbb{R})$, we may assume that $\psi \in \mathcal{F}(\mathbb{R})$. In fact, we may assume that $\psi = h_j$ for some $j \in \mathbb{N}$. Write $\psi = \sum_{k \in \mathbb{N}} c_k h_k$. For each $n \in \mathcal{N}$, write $\psi_n = \sum_k c_{n,k} \sigma_{n,k}$, where the index k runs over the range specified in Section 2. Now

$$c_j = \langle h_j | \psi \rangle = \lim_{n \in \mathcal{N}} \langle h_{n,j} | \psi_n \rangle = \lim_{n \in \mathcal{N}} c_{n,j}.$$

Putting $\epsilon := e^{2\pi i j t}$, then $\langle h_j | \hat{g}\psi \rangle = \epsilon c_j = \lim_{n \in \mathcal{N}} \epsilon c_{n,j} = \lim_{n \in \mathcal{N}} \langle h_{n,j} | \hat{g}_n \psi_n \rangle$. \square

References

1. G. G. Athanasiu, E. G. Floratos, Coherent states in finite quantum mechanics, *Nuclear Phys.* B 425 (1994), 343-364.
2. G. G. Athanasiu, E. G. Floratos, S. Nicolis, Holomorphic quantization on the torus and finite quantum mechanics, *J. Phys.* A 29 (1996), 6737-6745.
3. N. M. Atakishiev, S. K. Suslov, Difference analogues of the harmonic oscillator, (english translation) *Teoreticheskaya i Mat. Fiz.*, 85 (1990), 64-73.
4. R. Balian, C. Itzykson, Observations sur la mécanique quantique finie, *C. R. Acad. Sc. Paris, Ser. I*, 303 (1986), 773-778.
5. L. Barker, Continuum quantum systems as limits of discrete quantum systems, (Preprint).
6. L. Barker, Ç. Candan, T. Hakioglu, A. Kutay, H. M. Ozaktas, The discrete harmonic oscillator, Harper's equation, and the discrete fractional Fourier transform, *J. Phys. A: Math. Gen.* (to appear).
7. Ç. Candan, *The Discrete Fractional Fourier Transform* (Bilkent University M.S. thesis, July 1998, unpublished).
8. Ç. Candan, M. A. Kutay, H. M. Ozaktas, The discrete fractional Fourier transform, (pages 1713-1716 in) *Proceedings of the 1999 IEEE International Conference on Acoustics, Speech, and Signal Processing, Vol. III* (IEEE, Piscataway, New Jersey 1999).
9. T. Hakioglu, Linear canonical transformations and quantum phase: a unified canonical and algebraic approach, *J. Phys. A: Math. Gen.* 32 (1999), 4111-4130.
10. A. C. McBride, F. H. Kerr, On Namias' fractional Fourier transform, *IMA J. of Applied Math.* 39 (1987), 159-175.
11. J. H. McClellan, T. W. Parks, Eigenvalue and eigenvector decomposition of the discrete Fourier transform, *IEEE Trans. Audio Electroacoust.* AU-20 (1972), 66-74.
12. V. Namias, The fractional order Fourier transform and its application to quantum mechanics, *J. Inst. Math. Applics.* 25 (1980), 241-265.
13. H. M. Ozaktas, M. A. Kutay, D. Mendlovic, Introduction to the fractional Fourier transform and its applications (pages 239-291 in) P. W. Hawkes (editor) *Advances in Imaging and Electron Physics, Vol. 106*, (Academic Press, San Diego, California 1998).
14. H. M. Ozaktas, D. Mendlovic, M. A. Kutay, Z. Zalevsky, *The Fractional Fourier Transform: with Applications to Optics and Signal Processing* (Wiley, New York, to appear).

15. S-C. Pei, M-H. Yeh, Improved discrete fractional Fourier transform, *Optics Lett.* 22 (1997), 1047-1049.
16. R. Picard, *Hilbert space approach to some classical transforms*, (Pitman Research Notes in Math. 196, Longman, Harlow U.K., 1989).
17. R. Rammal, J. Bellisard, An algebraic semi-classical approach to Bloch electrons in a magnetic field, *J. Phys. France*, 51 (1990), 1803-1830.
18. J. H. Wilkinson, *The algebraic eigenvalue problem*, (Oxford Univ. Press, 1965).